



KIDS Clinical Guideline - Drug dose calculator

All drugs to be given by intravenous route unless otherwise stated. It is the responsibility of the clinician to ensure drugs are used appropriately according to the clinical situation and doses double checked. KIDS/NTS does not accept any liability.

Date Weight (kg): Name:

Weight checked by: Name..... Signed..... Prescriber : Name..... Signed.....

Single drug doses:	
Induction agents for intubation	Dose
Ketamine (2 mg/kg)	140mg max
Thiopentone (2-4 mg/kg)	160 - 280mg max
Fentanyl (1-5 micrograms/kg)	80 - 350micrograms max
Propofol (2-5 mg/kg)	160 - 350mg max
Muscle relaxants for intubation	
Suxamethonium (Weight <10kg: 2mg/kg)	-
Suxamethonium (Weight >=10kg: 1mg/kg)	80mg
Rocuronium (1mg/kg)	80mg
Vecuronium (0.1mg/kg)	8 mg
Emergency drugs for cardiac arrest	
Adrenaline 1:10,000 (0.1ml/kg)	8ml
Atropine (20micrograms/kg)	1mg max
Adenosine (0.1 - 0.5 mg/kg)	6-12mg
Amiodarone (5mg/kg)	300 mg max
Calcium gluconate 10% (0.5ml/kg)	20ml max
Sodium Bicarbonate 8.4% (1ml/kg)	50ml max
Anticonvulsants and drugs for raised ICP	
Lorazepam (0.1mg/kg)	4mg max
Phenytoin (20mg/kg - over 20 mins)	1600mg
Phenobarbitone (20mg/kg)	1g max
Paraldehyde 50%:Olive oil 50% (0.8ml/kg)	20ml PR max
Mannitol dose (0.25 - 0.5 g/kg)	20 - 40grams
Mannitol volume Peripheral: 10% solution	200 - 400 ml
Mannitol volume Central: 20% solution	100 - 200 ml
3% saline (3ml/kg)	240ml

Drug doses for infusion:			Total		
Sedation	Amount	Diluent - see key*	Volume	Rate	Dose
Morphine	80mg	D5/D10/NS	50ml	0.5 - 2 ml/hr	10-40micrograms/kg/h
Midazolam	100mg	D5/D10/DS/NS	50ml	1.2 - 7.2 ml/hr	0.5-3 micrograms/kg/min
Paralysis					
Rocuronium	Neat solution 300mg in 30ml			4.8 - 8 ml/hr	0.6 - 1mg/kg/h
Vecuronium	480mg	D5/D10/NS	50ml	0.5 - 2ml/hr	1 - 4micrograms/kg/min
Vasoactive drugs					
Dopamine	<i>Peripheral</i> 80mg	D5/D10/NS	50ml	15.0 - 60.0ml/hr	5 - 20micrograms/kg/min
	<i>Central</i> 625mg max	D5/D10/NS	50ml	1.9 - 7.7 ml/hr	5 - 20micrograms/kg/min
Dobutamine	<i>Peripheral</i> 80mg	D5/D10/NS	50ml	15.0 - 60.0ml/hr	5 - 20micrograms/kg/min
	<i>Central</i> 625mg max	D5/D10/NS	50ml	1.9 - 7.7 ml/hr	5 - 20micrograms/kg/min
Adrenaline	<i>Peripheral</i> 1mg	D5/D10/NS	50ml	2.4-120ml/hr	0.01 - 0.5 micrograms/kg/min
	<i>Central</i> 8mg	D5/D10/NS	50ml	0.3-15ml/hr	0.01 - 0.5micrograms/kg/min
Noradrenaline	<i>Central</i> 8mg	D5/DS	50ml	0.3-15ml/hr	0.01 -0.5micrograms/kg/min
Ductal patency drugs omitted as patient over 10kg					
-	-	-	-	-	-
-	-	-	-	-	-
Asthma drugs					
Salbutamol	For Salbutamol bolus dose please refer to BNFC				
	<i>Infusion Peripheral</i> 10mg	D5/NS	50ml	24.0 - 48.0 ml/hr	1 - 2micrograms/kg/min
	<i>Infusion Central</i> 25mg	D5/NS	50ml	9.6 - 19.2 ml/hr	1 - 2micrograms/kg/min
Caution - monitor for toxicity if salbutamol dose higher than 20mcg/min					
Aminophylline	Loading dose:		400mg over 20 minutes; use 500mg/500ml concentration as per infusion		
Use ideal wt if obese	500 mg	D5	500ml	80 ml/hr	1mg/kg/h
If patient is aged greater than 12 years please refer to BNFC for aminophylline dosing					
Magnesium sulphate 50%	4ml	D5/NS	100 ml	300 ml/hr	2 grams over 20 mins
DKA drugs					
Insulin	50units	D5/D10/NS	50ml	8.0ml/hr	0.1units/kg/h

*Diluent key: D5 = 5% glucose D10 = 10% glucose NS = 0.9% saline DS = 0.45% saline and 5% glucose